



NPK-fertilizer solution Nordfert 8-8-8 + microelements

6.4-6.4-6.4 + Microelements

Mass percent composition (%):

Total nitrogen (N)	6,4
Ammoniacal nitrogen (NH ₄ -N)	3,87
Nitric nitrogen (NO ₃ -N)	2,55
Total phosphorus pentoxide (P ₂ O ₅), water soluble	6,4
Total potassium oxide (K ₂ O), water soluble	6,4

Microelements

Boron (B)	0,015
Copper (Cu)	0,03
Iron (Fe)	0,06
Manganese (Mn)	0,03
Molybdenum (Mo)	0,005
Zinc (Zn)	0,01

Density: 1.25 g/ml

8-8-8+microelements

Volume percent composition (%):

Total nitrogen (N)	8
Ammoniacal nitrogen (NH ₄ -N)	4,84
Nitric nitrogen (NO ₃ -N)	3,19
Total phosphorus pentoxide (P ₂ O ₅), water soluble	8
Total potassium oxide (K ₂ O), water soluble	8

Microelements

Boron (B)	0,018
Copper (Cu)	0,035
Iron (Fe)	0,07
Manganese (Mn)	0,035
Molybdenum (Mo)	0,006
Zinc (Zn)	0,012

Date of manufacture: 00.00.0000

Expiration date: 00.00.0000

Registration number: 00000000

Safety and storage measures

To maintain the safety of the product quality packaging of fertilizers should be stored in a cool, dry, indoor place at temperature between +10 and +25°C, apart from the food and drugs. Avoid storage near flammable gases, electric motors and internal combustion engines, instruments, and chemicals. Do not leave empty packages in places easily accessible to children or pets. The usual precautions for handling fertilizers should be observed. Wash hands and face after handling the product. In case of contact with eyes, flush with water. After contact with skin, wash with water and soap if necessary. If ingested, rinse mouth and drink plenty of water. Take steps to control product discharge into the environment. Notify appropriate authorities if required and arrange for the recovery of spilled material. Methods for Clean-up: Dilute and wash away with large amounts of water. Dispose of in accordance with local regulation.



Advantages:

- Fully water soluble balanced fertilizer enriched with micronutrients.
- Activates enzymes and promotes active and healthy development.
- Provides better cell division, cell enlargement, and strengthens cell walls.
- Enhances seed germination, durability, and protein content.
- Contributes to improved flowering, fruit set and fruiting.
- Helps improve the quality of crops and increases yields.
- Improves the taste, color and shape of fruit.
- Maintains nutrient balance.
- Increases ability to tolerate cold, diseases, insects and pests.

Application:

Can be used in all fertigation systems and foliar application. Can be mixed with plant protection. A preliminary mixing of fertilizer with plant protection in a small cup is recommended. Shake well before use.

Crop	Growth stage	Dosage (l/ha)
Grains	Tillering	4-6
	Stem elongation	
	Heading	4-6
	After flowering	
Oilseed rape, linen	Beginning of vegetation	4-7
	10-14 days after	
	Phase of green bud	
Sugar and fodder beet	4-10 leaf phase	4-6
	10-14 days after	
	Before closing within the rows	5-7
Potatoes	Complete germination	4-8
	Before flowering	
	After flowering	
Corn	6-8 leaf phase	4-6
	Every 10-14 days - 2 applications	
Legumes	Before flowering	5-8
	After flowering	
	7-10 days after	
Strawberry	The end of flowering	4-6
	Beginning of berry growth	
	7-10 days later	
Fruit trees and shrubs	The end of flowering	6-9
	After flowering	
	10-14 days later	
Root vegetables	2-3 weeks after shooting	5-6
	Every 10-14 days - 2 applications	
Vegetables under cover: tomato, pepper, cucumber, lettuce, radish, celery, etc.	3 leaf phase - formation of germs	0.5-1 l per 100 l of water
	2-3 sprayings every 10-14 days	0.75-1 l per 100 l of water
Ornamentals and others	0.5-0.75 l of fertilizer per 100 l of solution	
Recommended amount of water: field crops - 150-300 l/ha; greenhouse and garden - 150-300 l/ha		

Net: 20 l (25 kg)

Country of origin: Estonia / EU
Producer / Supplier: NordFert OÜ
Fosforiidi 4, Maardu, 74114, Estonia
Email: info@nordfert.com