



Iron fertilizer solution

NordFert Micro Fe 100 g/l

Mass percent composition (%):

Iron (Fe), water soluble	7,5
--------------------------	-----

Density: 1.35 g/ml

Volume percent composition (%):

Iron (Fe), water soluble	10,0
--------------------------	------

Date of manufacture: 00.00.0000
Expiration date: 00.00.0000
Registration number: 000000000

Safety and storage measures

To maintain the safety of the product quality packaging of fertilizers should be stored in a cool, dry, indoor place at temperature between +10 and +25°C, apart from the food and drugs. Avoid storage near flammable gases, electric motors and internal combustion engines, instruments, and chemicals. Do not leave empty packages in places easily accessible to children or pets. The usual precautions for handling fertilizers should be observed. Wash hands and face after handling the product. In case of contact with eyes, flush with water. After contact with skin, wash with water and soap if necessary. If ingested, rinse mouth and drink plenty of water. Take steps to control product discharge into the environment. Notify appropriate authorities if required and arrange for the recovery of spilled material. Methods for Clean-up: Dilute and wash away with large amounts of water. Dispose of in accordance with local regulation.



Advantages:

- Promotes formation of chlorophyll and its functioning.
- Improves nitrogen uptake.
- Stimulates proper growth and cell division.
- Increases formation of enzymes and proteins.
- Improves the process of respiration and metabolism.
- Can be used for all types of fruits and plants throughout different growth stages.

Application:

Can be used in all fertigation systems and foliar application. Especially recommended in case of weak development or nutrient deficiency symptoms. Use only in case of need. Do not exceed recommended dosages. In case of use with plant protection a preliminary mixing is recommended. Shake gallon well before use.

Crop	Growth stage	Dosage (l/ha)
Grains	Beginning of tillering; Tillering; Stem elongation; Beginning of head formation.	0.5-1.5
Oilseed rape	Rosette formation; Beginning of vegetation; Beginning of stem elongation; Green bud phase; Beginning of flowering.	0.5-1.5
Corn	3-4 leaf stage; 6-10 leaf stage; Before flowering.	1-2
Legumes	Vegetative growth; Before flowering; After flowering.	1-2
Potato	Complete shooting; Before flowering; Formation of green tubers.	1-2
Beet	4-10 developed leaves; 10-14 days later; Before closing within the rows.	1-2
Recommended amount of water: field crops - - 150-300 l/ha; greenhouse and garden - 150-300 l/ha		

Net: 20 l (27 kg)

Country of origin: Estonia / EU
Producer / Supplier: NordFert OÜ
Fosforiidi 4, Maardu, 74114, Estonia
Email: info@nordfert.com